

Presynaptic calcium dynamics of neurons in answer to various stimulation protocols

Michael Meyer-Hermann, Frido Erler, Gerhard Soff

Institut für Theoretische Physik, TU Dresden, D-01062 Dresden, Germany
E-Mail: meyer-hermann@physik.tu-dresden.de

Abstract: We present a new model for the dynamics of the presynaptic intracellular calcium concentration in neurons evoked by various stimulation protocols. The aim of the model is twofold: We want to discuss the calcium transients during and after specific stimulation protocols as they are used to induce long-term-depression and long-term-potentialiation. In addition we would like to provide a general tool which allows the comparison of different calcium experiments. This may help to draw conclusions on a wider base in future.

Introduction

A most fascinating challenge in neuron biology is a deep understanding of the mechanisms involved in long-term-effects (LTE) such as long-term-potentialiation (LTP) and long-term-depression (LTD). The multiplicity of possibly important mechanisms is immense and is explored in a great variety of experiments [1]. However, the interpretations of those experiments are not as conclusive as they could be if it was possible to better compare experiments executed on different systems. We claim that a lot of detailed information on LTE is hidden in presently available experiments.

One possible way to uncover this hidden knowledge is to construct a tool which is able to translate different experiments into a common language and then to compare them quantitatively. Such a tool is provided here for the analysis of the intracellular calcium concentration in presynaptic nerve terminals. It is very well established by experiments that LTE are mostly connected with dynamical changes in the calcium concentration [2]. Larger calcium concentrations induce a lot of mechanisms that may influence the transmission efficiency of synapses. This applies not only to the postsynaptic side of the synapse, where an calcium influx is believed to be necessary for the induction of LTP [3]. Intracellular calcium also induces exocytosis on the presynaptic side of the synapse and therefore is in parts responsible for the release of neurotransmitter into the synaptic cleft [4]. It is believed that calcium/calmodulin dependent protein kinases may trigger the amount of released neurotransmitter. This has been shown for example for the calcium/calmodulin kinase II in the squid giant synapse [5]. Therefore, we think that the intracellular calcium concentration is an appropriate observable to study the induction of effects as LTD and LTP from the very beginning.

Already on this early level of LTE-induction a comparative quantitative evaluation of different experiments may lead to new insights. Corresponding experiments

have been done using different systems or using the same system under different conditions. In addition, a dependence of the presynaptic calcium dynamics on the corresponding target cell has been observed [6]. In order to compare those experiments quantitatively one has to determine characteristics of the experiment that are important for the calcium dynamics and to introduce them into the model terminology.

In the following we will develop a corresponding model which is based on previously developed models of the intracellular calcium dynamics. The number of models describing presynaptic calcium dynamics is comparably small [7, 8, 9, 10]. The main focus of model work in this field lies on the postsynaptic calcium dynamics [7, 11, 12, 13, 14, 15, 16, 17, 18] (for a review we refer to [19]). This is surely related to the fact that dendrites are better accessible in experiment than presynaptic boutons. Therefore, we use the knowledge which has been established on the postsynaptic side of the synapse in order to build a new model which can be used for comparison of presynaptic calcium dynamics in different experiments. Presynaptic measurements became more frequent due to technical developments in the last years [20].

The model has to be adjusted to specific experiments in a well-defined procedure, which is illustrated for the example of presynaptic nerve terminals in the rat neocortex [6]. The results found here are discussed in the context of LTE. For more details of the model and the results presented here we refer to [21]. This especially concerns a more detailed analysis of the model assumptions and the robustness of the results.

The model on the level of single proteins

We construct a new deterministic one-compartment model for the presynaptic calcium dynamics. The intracellular calcium concentration is changed in response to variations of the membrane potential by an ion-flux through voltage-gated calcium channels. The flux is driven by the electro-chemical gradient at the membrane. Intracellular calcium is bound by an endogenous buffer. Only a small number of ions remains free and increases the concentration of free calcium. An increased calcium concentration is reduced by two types of transport proteins: ATP-driven calcium pumps and natrium-calcium exchangers. Their activity basically depends on the free calcium concentration. A leakage transmembrane calcium current is assumed, that compensates the activity of transport proteins and channels in the rest state of the cell.

The model is formulated in terms of a set of ordinary coupled differential equations for the intracellular calcium concentration c :

$$\frac{dc}{dt} = \frac{G}{zF} \frac{\rho_U J_U (g_U(U), U(t), \overline{U}(c)) - \rho_P J_P(g_P(c)) - \rho_E J_E(g_E(c)) + L}{1 + \Theta_b(c) + \Theta_i(c)} \quad (1)$$

where $\rho_{U,P,E}$ are the surface densities of the voltage-gated channels (HVA), the PMCA-type calcium pumps, and the natrium-calcium exchanger (type 1), respectively.

$$J_{U,P,E} = g_{U,P,E} I_{U,P,E} \quad (2)$$

are the corresponding single protein currents. They are determined by the single protein open probabilities $g_{U,P,E}$, where $g_{P,E}$ depend on c according to standard Hill equations [22] for the pumps and the exchangers with Hill coefficient 2 [23] and 1 [24], respectively. The open probability for the voltage gated calcium channels (HVA) obeys [25]

$$\frac{dg_U(U)}{dt} = \frac{1}{\tau} \left\{ \left[1 + \exp \left\{ \frac{U_{1/2} - U}{\kappa} \right\} \right]^{-1} - g_U(U) \right\} , \quad (3)$$

with $\tau = 1ms$ the channel mean open time, $U_{1/2} = 3mV$ the half activation voltage, and $\kappa = 8mV$ the steepness factor. The values are taken from a representative HVA-channel [25]. I_U is the voltage dependent current which is assumed to follow Ohms law with open channel conductivity of $17pS$. The electrochemical gradient is calculated with respect to $\bar{U}(c)$, the (calcium dependent) calcium reversal potential, which depends on c according to the Nernst equation.

$I_P = 10^{-17}A$ and $I_E = 40 \cdot 10^{-17}A$ are the maximum activities of the transport proteins. G is a geometry factor (ratio of surface and volume in the compartment), $z = 2$ is the valence of the calcium ions, F the Faraday constant, and L the leak current which is determined by the steady state condition.

$U(t)$ is the stimulating transmembrane voltage function. We assume that the feedback effect of the calcium influx on the membrane potential is small with respect to standard action potentials. One may think of a voltage clamped situation. However, for large calcium concentrations as they may appear in high frequency stimulations, calcium buffers may saturate and the impact on the membrane potential may become non-negligible. Such effects are not considered in the present analysis.

The buffers (endogenous and indicator) are treated in a quasi-steady state approximation, which claims that the calcium binds and dissociates faster than the typical time scale under consideration. Then the dynamical behavior of the buffers reduces to a correction factor in Eq. (1) which depends on the calcium concentration only:

$$\Theta_b(c) = \frac{b_{\max} K_b}{(K_b + c)^2} \quad \text{and} \quad \Theta_i(c) = \frac{i_{\max} K_i}{(K_i + c)^2} . \quad (4)$$

Here $b_{\max}$ and $i_{\max}$ are the total concentrations of the endogenous buffer and the indicator, respectively. $K_{b,i}$ are the corresponding dissociation constants. Note that it is important to include the indicator used in the experiment, because the indicator is basically acting as additional buffer and may drastically change the amount of free intracellular calcium.

The differential equations are solved numerically and the solution describes the time course of the calcium concentration resulting in response to single action potentials or to series of action potentials (as they are used to induce LTD or LTP). The above mentioned aspired generality of the model is reflected in a separation of the model parameters into three classes, described in the following subsections.

Universality

The model is universal enough to be applicable to a wide class of different neuron

types. To this end the model is based on the experimental knowledge about *single* proteins which are postulated to have neuron-type independent properties. These are the single protein characteristics g_x (including all derived physiological properties) and the single protein transmembrane currents J_x , where x stands for the corresponding type of protein (see e.g. [23, 24, 25]).

Type specificity

The model is specific enough to be applicable to well defined neuron types. This is achieved by the introduction of measurable neuron-type specific parameters which has to be determined for each experiment separately. Basically, these are the protein densities ρ_x in the membrane. As no space resolution of the calcium concentration is considered, these densities may be thought as average values over the whole synaptic membrane. Also the concentration of the endogenous buffer $b_{\max}$ and its dissociation constant K_b belong to the neuron-type specific class of parameters. Finally, the surface to volume ratio G of the synaptic compartment quantitatively determine the concentration changes due to transmembrane currents.

Condition specificity

The model includes enough general specifications in order to adjust the model to specific experimental conditions. The form and amplitude of the action potential $U(t)$ is simulated with a system of coupled differential equations (not shown here) and can be adapted to the specific action potential used in experiment. The LTE-stimulation protocols used in experiment are simulated with a corresponding series of single action potentials. Intracellular calcium concentrations are generally visualized with the help of calcium indicators. They act as an additional buffer in the cell and, therefore, may influence the calcium dynamics. In the model they are treated in complete analogy to the endogenous buffer and are characterized by the indicator specific dissociation constant K_i and the used indicator concentration $i_{\max}$.

Adjustment to a specific experiment

The idea of this semi-universal model for presynaptic calcium dynamics is to determine the universal parameters using single protein experimental data and to maintain the resulting values for the evaluation of different systems and experimental conditions. *Universality* has to be understood as the postulated statement, that universal parameters are not the parameters that are most sensitive to a transfer from one experiment to another. The adjustment of the model to a specific experiment is achieved through the determination of the system-specific and condition-specific parameters. Note, that without any exception these are parameters with direct physiological interpretation. Therefore, most of them may be accessible up to a sufficient precision in several experiments.

In the described procedure the main part of the model is determined by sources that are independent of the experiment under consideration. Therefore, the value

of the model is tested by its capability of reproducing the answer of the calcium concentration to single action potentials on the basis of those independent and fixed universal parameters. As in general not all *specific* parameters will have been determined in the experiment under consideration, we will fit the remaining unknown parameters to the single action potential calcium response. With the help of a thus defined model it should be possible to analyse the measured calcium transients evoked by LTD or LTP stimulation protocols.

This has been executed for an experiment on single nerve terminals of pyramidal neurons in the neocortex of rats [6]: Most specific parameters used in the model are directly accessible in this experiment. This concerns for example the form of the action potential applied to the nerve terminal, the geometry (i.e. the surface to volume ratio $G = 3/\mu m$), the concentration of endogenous buffer $b_{\max} = 120\mu M$, and the characteristics and concentrations of the used calcium indicator (magnesium green). Therefore, we are in the situation that the whole model is a priori determined either by independent sources (concerning universal parameters) or by the available data from the experiment under consideration. The only unknown parameters are the surface densities of the calcium transport proteins. These are fitted to the measured calcium transient evoked by single action potentials ($\rho_U = 2.65/\mu m^2$, $\rho_P = 4000/\mu m^2$, and $\rho_E = \rho_P/30$). The result Fig. 1 shows, that the measured calcium transient is reproduced correctly.




Figure 1: The calcium transients evoked by single action potentials in single boutons of pyramidal neurons in the rat neocortex ($100\mu M$ magnesium green). The dotted line shows the experimental values [6] (we thank H.J. Koester and B. Sakmann for kind permission to reproduce the data), and the full line shows the model result with fitted protein densities.

LTE stimulation

In order to check if the thus defined model has predictive power, we calculate the intracellular calcium transients evoked by series of action potentials with varying frequency. Basically, the model parameters remain unchanged. As in the corresponding experimental setup (see [6] Fig.9) a different action potential (compared to Fig. 1) has been used, the action potential is adapted in the model and the channel densities are fitted to the *single* action potential calcium response, correspondingly ($\rho_U = 2.77/\mu m^2$, $\rho_P = 3400/\mu m^2$, and $\rho_E = \rho_P/30$). In addition the indicator concentration is increased, which is considerably higher in this experimental setup. Now the 10 Hz stimulus is applied and the result is shown in Fig. 2. The model




Figure 2: The calcium transients evoked by a 10 Hz stimulus in single boutons of pyramidal neurons in the rat neocortex ($500\mu M$ magnesium green). The full line represents the model result and the dotted line the corresponding measured transients [6] (we thank H.J. Koester and B. Sakmann for kind permission to reproduce the data). The dashed line represents the best single exponential fit to the experimental base line values [6].

result is in quantitative agreement with the calcium transients seen in the experiment: The intracellular calcium concentration reaches a new baseline level during the stimulation process which breaks down when the stimulus is switched off. The calcium concentration oscillates on the top of the new base line in coherence with the stimulation potential. The calcium reducing processes are slightly too slow in the model.

On this basis we can calculate the calcium transients in response to stimulation protocols with various frequencies. We find that calcium transients do not overlap

for low frequencies typical for LTD-induction. The emergence of a new baseline in the calcium concentration at frequencies around 10 Hz may be interpreted as threshold for the induction of LTP. Note, that this threshold frequency strongly depends on the used calcium indicator concentration. This is especially relevant for the interpretation of experimental results. A stimulation with frequencies around 50 Hz (typical for the induction of LTP) leads to a more pronounced enhancement of the calcium baseline. This qualitative behavior is in agreement with experiments carried out on dendritic spines of pyramidal neurons [26].

Conclusion

Our new model for transients of the presynaptic intracellular calcium concentration evoked by various stimulation protocols reproduces the general behavior observed in experiment. It is exclusively constructed with parameters that have a direct physiological interpretation. Its basis are the single proteins properties. The characteristics of single proteins are considered to be universal in the sense that they remain unchanged for different experimental setups. The model has been adjusted to a specific experiment that measured intracellular calcium transients in nerve terminals of pyramidal neurons of the rat neocortex. To this end the parameters specific for this experiment have been extracted from it or, if not available, have been fitted to the single action potential calcium response. The model results turned out to be in quantitative agreement with the experiment. This applies not only to the presynaptic calcium concentration response to single action potentials but also to 10 Hz stimuli. We did not find any reason for the involvement of calcium-induced-calcium-release in the induction of LTE. However, it seems that an additional mechanism (e.g. calcium channel inactivation) may be necessary to understand the induction of LTD on the level of presynaptic calcium transients.

More generally, the separation of universal and specific parameters enables us to analyse different results observed in several experiments. With the help of the new model one may decide, if those differences are significant or due to different experimental setups.

References

- [1] BAUDRY, M. & DAVIS, J.L. (EDS.), Long-Term Potentiation. *The MIT Press, Cambridge Massachusetts, London*. Vol. 1 (1991), Vol. 2 (1994), Vol. 3 (1996).
- [2] BLISS, T.V.P. & COLLINGRIDGE, G.L., A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* **361**, 31-39 (1993).
- [3] TSIEN, R.W. & MALINOW, R., Long-term potentiation: Presynaptic enhancement following postsynaptic activation of Ca^{++} -dependent protein kinases. *Cold Spring Harbor Symp. Quant. Biol.* **55**, 147-159 (1990).

- [4] DODGE, F.A. & RAHAMIMOFF, R., Co-operative action of calcium ions in transmitter release at the neuromuscular junction. *J. Physiol.* **193**, 419-432 (1967).
- [5] LLINAS, R., MCGUENNESS, T.L., LEONARD, C.S., SUGIMORI, M. & GREENGARD, P., Intraterminal injection of synapsin I or calcium/calmodulin-dependent protein kinase II alters neurotransmitter release at the squid giant synapse. *Proc. Natl. Acad. Sci. USA* **82**, 3035-3039 (1985).
- [6] KOESTER, H.J. & SAKMANN, B., Calcium dynamics associated with action potentials in single nerve terminals of pyramidal cells in layer 2/3 of the young rat neocortex. *J. Physiol.* **529.3**, 625-646 (2000).
- [7] NEHER, E. & AUGUSTINE, G.J., Calcium gradients and buffers in bovine chromaffin cells. *J. Physiol.* **450**, 273-301 (1992).
- [8] DE SCHUTTER, E., ANGSTADT, J.D. & CALABRESE, R.L. A model of graded synaptic transmission for use in dynamic network simulations. *J. Neurophysiol.* **69** 1225-1235 (1993).
- [9] HELMCHEN, F., BORST, J.G.G. & SAKMANN, B., Calcium dynamics associated with a single action potential in a CNS presynaptic terminal. *Biophys. J.* **72**, 1458-1471 (1997).
- [10] SINHA, S.R., WU, L.-G. & SAGGAU, P., Presynaptic calcium-dynamics and transmitter release evoked by single action potentials at mammalian central synapses. *Biophys. J.* **72**, 637-651 (1997).
- [11] GAMBLE, E. & KOCH, C., The dynamics of free calcium in dendritic spines in response to repetitive synaptic input. *Science* **236**, 1311-1315 (1987).
- [12] HOLMES, W.R. & LEVY, W.B., Insights into associative long-term potentiation from computational models of NMDA receptor-mediated calcium influx and intracellular calcium concentration changes. *J. Neurophysiol.* **63**, 1148-1168 (1990).
- [13] JAFFE, D.B., ROSS, W.N., LISMAN, J.E., LASSER-ROSS, N., MIYAKAWA, H. & JOHNSTON, D., A model for dendritic Ca^{2+} accumulation in hippocampal pyramidal neurons based on fluorescence imaging measurements. *J. Neurophysiol.* **71**, 1065-1077 (1994).
- [14] SIEGEL, M., MARDER, E. & ABBOTT, L.F., Activity-dependent current distributions in model neurons. *Proc. Natl. Acad. Sci. USA* **91**, 11308-11312 (1994).
- [15] SCHIEGG, A., GERSTNER, W., RITZ, R. & VAN HEMMEN, J.L., Intracellular Ca^{2+} stores can account for the time course of LTP induction: A model of Ca^{2+} dynamics in dendritic spines. *J. Neurophysiol.* **74**, 1046-1055 (1995).

- [16] DE SCHUTTER, E., Dendritic Voltage and Calcium-Gated Channels Amplify the Variability of Postsynaptic responses in a Purkinje-Cell Model. *J. Neurophysiol.* **80**, 504-519 (1998).
- [17] VOLFOVSKY, N., PARNAS, H., SEGAL, M. & KORKOTIAN, E., Geometry of Dendritic Spines Affects Calcium Dynamics in Hippocampal Neurons: Theory and Experiments. *J. Neurophysiol.* **82**, 450-462 (1999).
- [18] HOLMES, W.R., Models of calmodulin trapping and CaM kinase II activation in a dendritic spine. *J. Comput. Neurosci.* **8**, 65-85 (2000).
- [19] DE SCHUTTER, E. & SMOLEN, P., Calcium dynamics in large neuronal models, in: Koch, C. & Segev, I. (eds.), *Methods in neuronal modeling*. The MIT Press, Cambridge (Mass.), London 1998.
- [20] SABATINI, B.L. & REGEHR, W.G., Optical Measurement of Presynaptic Calcium Currents. *Biophys. J.* **74**, 1549-1563 (1998).
- [21] ERLER, F., MEYER-HERMANN, M. & SOFF, G., A quantitative model for the presynaptic calcium dynamics during the induction of long-term effects in neurons. *In preparation* (2002).
- [22] ZADOR, A., KOCH, C. & BROWN, T.H. Biophysical model of a Hebbian synapse. *Proc. Natl. Acad. Sci. USA* **87**, 6718-6722 (1990).
- [23] ELWESS, N.L., FILOTEO, A.G., ENYEDI, A. & PENNISTON, J.T., Plasma Membrane Ca^{2+} Pump Isoforms 2a and 2b are Unusually Responsive to Calmodulin and Ca^{2+} . *J. Biol. Chem.* **272**, 17981-17986 (1997).
- [24] BLAUSTEIN, M.P. & LEDERER, W.J., Sodium/Calcium Exchange: Its Physiological Implications. *Physiol. Rev.* **79**, 763-854 (1999).
- [25] MAGEE, J.C. & JOHNSTON, D., Characterization of single voltage-gated Na^{+} and Ca^{2+} channels in apical dendrites of rat CA1 pyramidal neurons. *J. Physiol.* **487.1**, 67-90 (1995).
- [26] HELMCHEN, F., IMOTO, K. & SAKMANN, B., Ca^{2+} Buffering and Action Potential-Evoked Ca^{2+} Signaling in Dendrites of Pyramidal Neurons. *Biophys. J.* **70**, 1069-1081 (1996).